



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2016

Matrilineal inheritance of a key mediator of prenatal maternal effects

Tschirren, Barbara ; Ziegler, Ann-Kathrin ; Pick, Joel L ; Okuliarová, Monika ; Zeman, Michal ;
Giraudeau, Mathieu

Abstract: Sex-linkage is predicted to evolve in response to sex-specific or sexually antagonistic selection. In line with this prediction, most sex-linked genes are associated with reproduction in the respective sex. In addition to traits directly involved in fertility and fecundity, mediators of maternal effects may be predisposed to evolve sex-linkage, because they indirectly affect female fitness through their effect on offspring phenotype. Here, we test for sex-linked inheritance of a key mediator of prenatal maternal effects in oviparous species, the transfer of maternally derived testosterone to the eggs. Consistent with maternal inheritance, we found that in Japanese quail (*Coturnix japonica*) granddaughters resemble their maternal (but not their paternal) grandmother in yolk testosterone deposition. This pattern of resemblance was not due to non-genetic priming effects of testosterone exposure during prenatal development, as an experimental manipulation of yolk testosterone levels did not affect the females' testosterone transfer to their own eggs later in life. Instead, W chromosome and/or mitochondrial variation may underlie the observed matrilineal inheritance pattern. Ultimately, the inheritance of mediators of maternal effects along the maternal line will allow for a fast and direct response to female-specific selection, thereby affecting the dynamics of evolutionary processes mediated by maternal effects.

DOI: <https://doi.org/10.1098/rspb.2016.1676>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-126758>

Journal Article

Accepted Version

Originally published at:

Tschirren, Barbara; Ziegler, Ann-Kathrin; Pick, Joel L; Okuliarová, Monika; Zeman, Michal; Giraudeau, Mathieu (2016). Matrilineal inheritance of a key mediator of prenatal maternal effects. *Proceedings of the Royal Society of London, Series B: Biological Sciences*, 283(1838):20161676.

DOI: <https://doi.org/10.1098/rspb.2016.1676>

1 **Matrilineal inheritance of a key mediator of prenatal maternal effects**

2

3 Barbara Tschirren^{1*}, Ann-Kathrin Ziegler¹, Joel L. Pick¹, Monika Okuliarová², Michal
4 Zeman² & Mathieu Giraudeau^{1, 3}

5

6 ¹Department of Evolutionary Biology and Environmental Studies, University of
7 Zurich, Winterthurerstrasse 190, 8057 Zurich, Switzerland

8

9 ²Department of Animal Physiology and Ethology, Faculty of Natural Sciences,
10 Comenius University, Bratislava, Slovak Republic

11

12 ³Centre for Ecology & Conservation, College of Life and Environmental Sciences,
13 University of Exeter, Penryn, UK

14

15

16 *Correspondence: Barbara Tschirren, Department of Evolutionary Biology and
17 Environmental Studies, University of Zurich, Winterthurerstrasse 190, CH-8057
18 Zurich, Switzerland; email: barbara.tschirren@ieu.uzh.ch, phone: +41 44 635 47 77,
19 Fax: +41 44 635 47 80

20

21 **Running title:** Inheritance of maternal effectors

22

Abstract

Sex-linkage is predicted to evolve in response to sex-specific or sexually antagonistic selection. In line with this prediction, most sex-linked genes are associated with reproduction in the respective sex. In addition to traits directly involved in fertility and fecundity, mediators of maternal effects may be predisposed to evolve sex-linkage because they indirectly affect female fitness through their effect on offspring phenotype. Here we test for sex-linked inheritance of a key mediator of prenatal maternal effects in oviparous species, the transfer of maternally-derived testosterone to the eggs. Consistent with maternal inheritance, we found that in Japanese quail (*Coturnix japonica*) granddaughters resemble their maternal, but not their paternal grandmother in yolk testosterone deposition. This pattern of resemblance was not due to non-genetic priming effects of testosterone exposure during prenatal development, as an experimental manipulation of yolk testosterone levels did not affect the females' testosterone transfer to their own eggs later in life. Instead, W chromosome and / or mitochondrial variation may underlie the observed matrilineal inheritance pattern. Ultimately, the inheritance of mediators of maternal effects along the maternal line will allow for a fast and direct response to female-specific selection, thereby affecting the dynamics of evolutionary processes mediated by maternal effects.

Keywords: maternal effects; yolk androgens; sex-specific selection; *Coturnix japonica*; hormones

44 **Introduction**

45 Sexual antagonism is common in nature and has important consequences for the
46 genomic arrangement of loci under sex-specific selection, as well as their inheritance
47 [1-3]. Indeed, because daughters are more likely to obtain high female-fitness alleles
48 from their mother than from their father, and vice versa, sex-specific (or sexually
49 antagonistic) selection will favour sex-linkage of traits differentially linked to male
50 and female fitness [4, 5]. A classic example for the evolution of sex-linkage in
51 response to sexually antagonistic selection is coloration in guppies (*Poecilia*
52 *reticulata*), which is associated with attractiveness in males [6], but makes males and
53 females more vulnerable to predation [7]. In response to these conflicting selection
54 pressures, a large proportion of the genetic variation in coloration has become linked
55 to the male-specific Y chromosome [8].

56 Even when selection is not acting in a sexually *antagonistic* way, sex-linkage may be
57 adaptive because it allows for a faster and more direct response to sex-specific
58 selection. Furthermore, if a trait is expressed in a sex-limited way, sex-linkage
59 prevents deleterious alleles from being sheltered from selection in the non-expressing
60 sex [4], again accelerating adaptive responses to selection. In line with these ideas,
61 male-specific fitness traits, such as sperm motility [9] or spermatogenesis [10], are
62 linked to the male-specific Y chromosome in species where the male is the
63 heterogametic sex (XY). And similarly, in species where the female is the
64 heterogametic sex (ZW), female fecundity and fertility traits are associated with the
65 female-specific W chromosome [11, 12].

66 We propose that in addition to traits directly involved in fecundity and fertility,
67 mediators of maternal effects (i.e. maternally-expressed traits that affect offspring
68 phenotype) may be predisposed to evolve sex-linkage because they indirectly affect

female fitness through their effect on offspring phenotype [13]. Furthermore, we argue that the potential for such sex-linkage of maternal effects mediators is particularly high in taxa where the female is the heterogametic sex (such as birds). Here we used a three-generation breeding design (ESM 1) in a captive Japanese quail (*Coturnix japonica*) population to test for sex-linkage of a key mediator of prenatal maternal effects in birds: the transfer of maternally-derived testosterone (T) to the eggs (yolk T transfer) [14-16]. Maternally transferred T affects a wide range of morphological, physiological, behavioural and life history traits in the offspring (i.e. it acts as a mediator of maternal effects [14-16]), and the costs and benefits of T exposure during prenatal development appear to depend on the social and environmental conditions encountered by the offspring [17-19]. Yolk T transfer is known to be heritable [20-22], but the design of previous studies did not allow to detect potential sex-linkage. We predict that if yolk T transfer is inherited along the maternal line, females will resemble their maternal, but not their paternal grandmother in their transfer of T to the eggs.

Material and methods

Study population

The study was conducted in a population of Japanese quail kept at the University of Zurich, Switzerland. Males and females were housed in separate outdoor aviaries (7 x 5.5 m each). For breeding, male-female pairs were transferred to cages (122 x 50 x 50 cm) within our facility. Cages contained *ad libitum* food, water, grit, a source of calcium, a shelter and a sand bath. The bottom of the cages was lined with sawdust. The breeding facility was kept on a 16 h : 8 h light : dark cycle at $20 \pm 3^{\circ}\text{C}$ (see [23] for a detailed description of animal husbandry).

94

95 *Egg collection, incubation and offspring rearing*

96 Eggs were collected daily, labelled with a non-toxic marker, and weighed. To
97 standardise incubation and rearing conditions, we artificially incubated the eggs
98 (mean \pm SD: 9.5 ± 0.84 eggs per female) (Favorit, HEKA Brutgeräte, Germany;
99 37.8°C , 55% humidity). For hatching, eggs were placed in individual containers to be
100 able to determine which chick hatched from which egg. After hatching, chicks were
101 raised in heated cages in mixed family groups (109 x 57 x 25 cm, Kükenaufzuchtbox
102 4002/C, HEKA Brutgeräte, Germany). Variation in the number of eggs laid while in
103 the breeding cages was small and there was no mother-daughter resemblance in the
104 number of eggs laid (generalised linear mixed model: $\chi^2 = 0.264$, $P = 0.607$).
105 For the yolk T analysis, yolk and albumen of one egg per female (the 5th) were
106 separated, weighed, homogenised, and frozen at -20°C . Previous work has shown that
107 within-clutch variation in yolk T concentration is small in Japanese quail (within-
108 female repeatability across different stages of the reproductive cycle > 0.7 [24]) and
109 the 5th egg is thus representative of a female's yolk T deposition to her eggs. Yolks
110 were collected across three generations (hereafter referred to as maternal and paternal
111 grandmothers, mothers, and (grand-) daughters) to assess the inheritance pattern (see
112 ESM 1). Within a generation, all females had the same age and had experienced the
113 same period of reproductive activity when eggs were collected.

114

115 *Yolk testosterone analysis*

116 Yolk T extraction and radioimmunoassay were performed following previously
117 published protocols [22]. In short, 100-110 mg of yolk were spiked with
118 approximately 2500 dpm of [^3H]-testosterone (PerkinElmer, USA) and extracted

twice with a mixture of diethyl and petroleum ether (7 : 3). Yolk T concentrations (pg / mg yolk) were quantified in 10 µl aliquots using [1,2,6,7-³H]-testosterone (PerkinElmer, USA, specific activity 63.47 Ci/mmoL) and a specific antibody generated in rabbits against testosterone-3-(carboxy-methyl) oxime bovine serum albumin conjugate [25]. The sensitivity of the assay was 1.62 ± 0.17 pg per tube. The mean recovery rate \pm SD was $79.3 \pm 6.4\%$. The samples were analysed in two assays. The intra- and inter-assay coefficients of variation were 4.7% and 6.5%, respectively. To test for (matrilineal) inheritance of yolk T transfer, we analysed the yolk T concentration in the eggs of 22 maternal grandmothers, 24 paternal grandmothers, 29 mothers and 40 (grand-) daughters (ESM 1). Yolk T concentrations were log transformed and standardised within generation before analysis to ensure normality of the residuals and equal variances across generations.

Yolk testosterone manipulation

To explore whether the resemblance in yolk T transfer along the maternal line (see Results) is due to non-genetic priming effects, we experimentally manipulated yolk T levels in eggs and tested 1) if T levels experienced during a female's prenatal development affect the transfer of T into her own eggs later in life, and 2) if the manipulation affects the transfer of T into the eggs of the daughters of these females (i.e. if the manipulation has a transgenerational effect). To this end, we experimentally increased yolk T concentrations in the eggs of half of the females of the second generation before incubation. This manipulation simulates an environmental effect on maternal yolk T transfer (i.e. an environmental maternal effect), as for example

observed in response to breeding density [26, 27], food availability [28, 29] or parasite abundance [19].

We injected eggs with 15 ng testosterone (Sigma-Aldrich, Switzerland) dissolved in 20 μ l safflower oil (Sigma-Aldrich, Switzerland) (T-treatment) or with 20 μ l safflower oil as a control (C-treatment). Clutches (N = 29) were assigned randomly to one of the two treatment groups. The injected dose is equivalent to approximately 1 SD of the yolk T content in the study population (mean \pm SD: 48.4 \pm 16.9 ng / yolk; range: 18.5 – 83.9 ng / yolk). Injections were performed at the pointed end of the egg, using an insulin syringe (Terumo, Belgium). The hole in the shell was closed with an adhesive film (Opsite, Smith & Nephew, Switzerland). There was no statistically significant difference in hatching success between T-injected and control eggs [30].

Furthermore, the yolk T manipulation did not significantly affect brood sex ratio (ESM 2). When females originating from T-manipulated and control eggs reached adulthood, we measured the T concentration they transferred to their own eggs (see above). Moreover, we measured the yolk T concentration in the eggs of 26 daughters of these females (as described above) to test for a transgenerational effect of the yolk T manipulation on yolk T transfer.

Statistical analysis

First, we used a linear mixed model to quantify the relationship between the yolk T concentration in the eggs of mothers (explanatory variable) and daughters (response variable). Family ID was included as a random effect to control for the non-independence of siblings.

Second, a similar model, this time with the T concentration in the eggs of the maternal and paternal grandmother as explanatory variables, was used in order to estimate the

relationship between the yolk T concentration in the eggs of both grandmothers and their granddaughters. To confirm the results of these linear mixed models, we conducted a model selection procedure using AICc criteria to determine if a model that contains maternal and / or paternal grandmother yolk T best explains yolk T transfer of granddaughters. Candidate models contained combinations of the maternal grandmother's and paternal grandmother's yolk T concentrations. All candidate models contained family ID as a random effect. Model selection was performed using the 'MuMIn' package [31] in R [32].

Third, we tested for an effect of the experimental yolk T manipulation on the transfer of yolk T later in life in 1) females that developed in the manipulated eggs (i.e. directly experienced manipulated T concentrations during their embryonic development), and 2) in the daughters of these females (to test for transgenerational effects of the manipulation) using linear mixed models that included T treatment, the yolk T concentration in the eggs of the mother and their interaction as fixed effects, and family ID as a random effect. For all linear mixed models, analyses were performed using the package 'lme4' [33] in R [32]. *P* values were obtained by comparing two nested models, with and without the variable of interest, using likelihood ratio tests.

Results

There was a significant positive relationship between the yolk T concentration in the eggs of mothers and daughters ($b \pm SE$: 0.437 ± 0.142 ; $\chi^2 = 8.185$, $P = 0.004$; Fig. 1A). Similarly, a significant positive relationship between the yolk T concentrations in the eggs of maternal grandmothers and granddaughters was found ($b \pm SE$: 0.366 ± 0.147 ; $\chi^2 = 5.415$, $P = 0.020$; Fig. 1B). In contrast, yolk T concentrations in the eggs of

paternal grandmothers and granddaughters were unrelated ($b \pm \text{SE}$: -0.027 ± 0.159 ; $\chi^2 = 0.001$, $P = 0.973$; Fig. 1C). In comparison, the resemblance in yolk mass between granddaughters and their maternal ($b \pm \text{SE}$: 0.266 ± 0.158) or paternal grandmother ($b \pm \text{SE}$: 0.250 ± 0.184) was very similar. As a consequence, analysing total yolk T content instead of yolk T concentration gave comparable results in all analyses. The finding that yolk T deposition is inherited along the maternal line was confirmed by a model selection procedure based on AICc, which revealed that a model containing only the maternal grandmother's yolk T concentration explained the granddaughters' yolk T transfer best. Models that contained additionally the paternal grandmother's yolk T concentration or only the paternal grandmother's yolk T concentration all had $\Delta\text{AICc} > 4.5$. There was no indication that an experimental increase of yolk T levels experienced during prenatal development influences a female's own transfer of yolk T later in life ($\chi^2 = 0.243$, $P = 0.622$; Fig. 2). Furthermore, the manipulation had no significant transgenerational effect on the yolk T transfer of the daughters of females that developed in the manipulated eggs ($\chi^2 = 0.035$, $P = 0.851$).

Discussion

Using a three-generation breeding design, we provide evidence for a significant within-family resemblance in the transfer of yolk T, an important mediator of prenatal maternal effects in oviparous species [15, 16]. However, in contrast to what is expected under autosomal inheritance, the resemblance in yolk T transfer between mothers and daughters, and between maternal grandmothers and granddaughters was very similar, whereas yolk T concentrations in eggs of paternal grandmothers and

217 granddaughters were unrelated. This pattern of resemblance is consistent with female-
218 linked inheritance.

219 Sex-linked inheritance can be caused by several non-mutually exclusive mechanisms.
220 First, information on the avian female-specific W chromosome, which is passed on
221 from mothers to daughters, may influence yolk T transfer. Although the W
222 chromosome contains only few genes [34, 35], it plays a key role in regulating female
223 fertility and fecundity [11, 12], likely through epistatic interactions between the W
224 chromosome and other parts of the genome [36]. Moreover, the expression of W
225 chromosome-linked genes has been found to rapidly respond to artificial selection on
226 female reproductive performance [12], again highlighting the important role of W-
227 linked variation in mediating female fitness.

228 Second, mitochondrial effects may underlie the observed maternal resemblance in
229 yolk T transfer. Mitochondria are, like W chromosomes, inherited along the maternal
230 line and there is accumulating evidence that mitochondrial genetic variation is non-
231 neutral [37, 38]. If mitochondrial variation affects yolk T transfer, for example by
232 influencing a female's metabolic rate [39], this could explain the female-linked
233 inheritance pattern. Indeed, there is a strong positive relationship between a female's
234 resting metabolic rate (RMR) and the amount of T she transfers to her eggs [40],
235 making this a plausible scenario. Interestingly, positive selection has shaped
236 *ATP5A1W*, a gene on the avian W chromosome that encodes a mitochondrial ATP
237 synthase subunit [41], suggesting that W- and mtDNA variation may epistatically
238 interact in shaping female-specific fitness traits [36]. Testing for associations
239 between sequence or structural [42] variation on the W-chromosome and / or the
240 mitochondria and variation in yolk T transfer will thus be a fruitful next step, and will

allow for an in-depth investigation of the molecular mechanisms underlying the maternal inheritance pattern observed in our study.

Besides sex-limited genetic variation, non-genetic mechanisms [43-45] may contribute to the resemblance in yolk T transfer along the maternal line. For example, prenatal exposure to yolk T may prime (‘program’) a female’s yolk T transfer to her own eggs at adulthood. Indeed, experimental manipulations have shown that variation in prenatal T exposure has long-term effects on both circulating T levels as well as T sensitivity later in life [46, 47]. We directly tested this hypothesis, but found no evidence that females originating from an egg with experimentally increased T concentration differed in their yolk T transfer from control females. Moreover, we found no evidence for a transgenerational effect of the yolk T manipulation on the deposition of yolk T in the next generation (i.e. in the daughters of females that developed in the manipulated eggs).

The former finding is in line with previous studies in pheasants (*Phasianus colchicus*)[48] and canaries (*Serinus canaria*)[49] that found no effect of experimentally increased prenatal T exposure on T transfer to the eggs. We can exclude that the lack of an effect was due to an unsuccessful manipulation, because the yolk T treatment affected a range of other behavioural and physiological traits in our study [30] as well as in [48] and [49]. Rather, it suggests that whereas prenatal exposure to T has long-term effects on both circulating T levels and T sensitivity [46, 47], it does not affect the transfer of T to the eggs.

Whereas we found no evidence that the T manipulation affected the (overall) yolk T transfer in the next two generations, the manipulation may differentially affect the deposition of yolk T to male and female eggs. However, this scenario appears unlikely given that evidence for differential allocation of T to male and female eggs is

266 weak across species [50], and absent in Japanese quail [51] (see also ESM 2).
267 Furthermore, although the T manipulation was performed within the natural range, it
268 is possible that the lack of a difference might be due to dose-response effects [52].
269 Given the highly controlled egg handling, incubation and chick rearing conditions in
270 our study, we can exclude that common postnatal environmental effects contribute to
271 the observed within-family resemblance. However as a third potential source of
272 matrilineal resemblance, other non-genetic effects such as the transmission of
273 epigenetic states across generations [45], other egg components (e.g. nutrients) that
274 indirectly prime yolk T transfer, or genomic imprinting may play a role. Although we
275 can currently not exclude such mechanisms, they are unlikely to explain our results
276 because to date neither the transgenerational transmission of epigenetic marks [53],
277 nor genomic imprinting [54, 55] have been documented in birds.
278 Ultimately, sex-linkage of yolk T transfer may have evolved in response to female-
279 specific selection and / or in order to resolve sexual conflict [3, 56]. Although yolk T
280 transfer is a trait that is expressed only in females, any underlying autosomal genes
281 might have pleiotropic effects on traits expressed in males as well [57]. For example,
282 yolk T transfer may not be independent of T levels in the circulation, on which strong
283 sexually antagonistic selection is acting on [58]. Interestingly, the relationship
284 between yolk T and plasma T levels differs across species [59], which may reflect
285 different stages in the resolution of this conflict. Under this scenario, we would
286 predict pronounced sex-linkage of yolk T transfer in species where yolk T and
287 circulating T levels are not correlated (anymore) (e.g. our study species [22]), but no
288 or limited sex-linkage in species where the two traits are (still) correlated (e.g. canary
289 *Serinus canaria* [60]).

In conclusion, we show that yolk T transfer, an important mediator of prenatal maternal effects in oviparous species, is inherited along the maternal line in Japanese quail. We can exclude the possibility that this maternal resemblance is due to common postnatal environmental effects or non-genetic priming effects of prenatal exposure to T on yolk T transfer later in life. Instead, our findings suggest that W-linked and / or mitochondrial variation might underlie the observed inheritance pattern. Female-linked inheritance of maternal effect mediators allows for a fast and direct response to female-specific selection and will thereby affect the dynamics of evolutionary processes mediated by maternal effects, such as the adaptation of populations to changing environments [61] or mother-offspring coadaptation [62].

Ethics

All procedures conform to the relevant regulatory standards and were conducted under licences provided by the Veterinary Office of the Canton of Zurich, Zurich, Switzerland (195/2010; 14/2014; 156).

Data accessibility

Data are available from Dryad (doi:10.5061/dryad.j76q1).

Competing interests

We have no competing interests.

312 **Authors' contributions**

313 BT conceived and coordinated the project, conducted the statistical analysis and wrote
314 the manuscript. AKZ, JLP and MG collected data and performed the egg
315 manipulation, AKZ, MO and MZ performed the hormone assays. All authors
316 commented on the manuscript.

317

318 **Acknowledgements**

319 We thank Alison Pick, Barbara Schnüriger, Pascale Hutter, Christina Ebner and Elif
320 Hanic for help with animal husbandry and data collection, and Erik Postma and two
321 anonymous reviewers for comments on the manuscript.

322

323 **Funding**

324 The study was financially supported by Schweizerischer Nationalfonds zur Förderung
325 der Wissenschaftlichen Forschung (PP00P3_128386 and PP00P3_157455) and Fonds
326 zur Förderung des akademischen Nachwuchses (FAN).

327

328 **Literature Cited**

- 329 1. Connallon, T. & Clark A. G. 2010 Sex linkage, sex-specific selection, and the
330 role of recombination in the evolution of sexually dimorphic gene expression
331 *Evolution* **64** (12), 3417-3442. (doi:10.1111/j.1558-5646.2010.01136.x).
- 332 2. Rice, W. R. 1984 Sex chromosomes and the evolution of sexual dimorphism
333 *Evolution* **38** (4), 735-742. (doi:10.2307/2408385).
- 334 3. Bonduriansky, R. & Chenoweth S. F. 2009 Intralocus sexual conflict. *Trends*
335 *Ecol. Evol.* **24** (5), 280-288. (doi:10.1016/j.tree.2008.12.005).
- 336 4. Day, T. & Bonduriansky R. 2004 Intralocus sexual conflict can drive the
337 evolution of genomic imprinting. *Genetics* **167** (4), 1537-1546.
338 (doi:10.1534/genetics.103.026211).
- 339 5. Mank, J. E. 2012 Small but mighty: the evolutionary dynamics of W and Y
340 sex chromosomes. *Chromosome Res.* **20** (1), 21-33. (doi:10.1007/s10577-011-9251-2).
- 341 6. Houde, A. E. 1997 *Sex, Color, and Mate Choice in Guppies*. Princeton, USA,
342 Princeton University Press.
- 343 7. Endler, J. A. 1980 Natural selection on color patterns in *Poecilia reticulata*.
344 *Evolution* **34** (1), 76-91.
- 345 8. Postma, E., Spyrou N., Rollins L. A. & Brooks R. C. 2011 Sex-dependent
346 selection differentially shapes genetic variation on and off the guppy Y chromosome.
347 *Evolution* **65** (8), 2145-2156. (doi:10.1111/j.1558-5646.2011.01314.x).
- 348 9. Chippindale, A. K. & Rice W. R. 2001 Y chromosome polymorphism is a
349 strong determinant of male fitness in *Drosophila melanogaster*. *Proc. Natl. Acad. Sci.*
350 *U S A* **98** (10), 5677-5682. (doi:10.1073/pnas.101456898).
- 351 10. Lange, J., Skaletsky H., van Daalen S. K. M., Embry S. L., Korver C. M.,
352 Brown L. G., Oates R. D., Silber S., Repping S. & Page D. C. 2009 Isodicentric Y

353 chromosomes and sex disorders as byproducts of homologous recombination that
 354 maintains palindromes. *Cell* **138** (5), 855-869. (doi:10.1016/j.cell.2009.07.042).

355 11. Hori, T., Asakawa S., Itoh Y., Shimizu N. & Mizuno S. 2000 Wpkci, encoding
 356 an altered form of PKCI, is conserved widely on the avian W chromosome and
 357 expressed in early female embryos: Implication of its role in female sex determination.
 358 *Mol. Biol. Cell* **11** (10), 3645-3660.

359 12. Moghadam, H. K., Pointer M. A., Wright A. E., Berlin S. & Mank J. E. 2012
 360 W chromosome expression responds to female-specific selection. *Proc. Natl. Acad.*
 361 *Sci. U S A* **109** (21), 8207-8211. (doi:10.1073/pnas.1202721109).

362 13. Mousseau, T. A. & Fox C. W. 1998 *Maternal Effects As Adaptations*. New
 363 York, Oxford University Press.

364 14. Schwabl, H. 1993 Yolk is a source of maternal testosterone for developing
 365 birds. *Proc. Natl. Acad. Sci. U S A* **90** (24), 11446-11450.

366 15. Groothuis, T. G. G., Müller W., von Engelhardt N., Carere C. & Eising C.
 367 2005 Maternal hormones as a tool to adjust offspring phenotype in avian species.
 368 *Neurosci. Biobehav. Rev.* **29**, 329-352.

369 16. Gil, D. 2008 Hormones in avian eggs: Physiology, ecology and behavior. *Adv.*
 370 *Study Behav.* **38**, 337-398. (doi:10.1016/s0065-3454(08)00007-7).

371 17. Muriel, J., Salmon P., Nunez-Buiza A., de Salas F., Perez-Rodriguez L.,
 372 Puerta M. & Gil D. 2015 Context-dependent effects of yolk androgens on nestling
 373 growth and immune function in a multibrooded passerine. *J. Evol. Biol.* **28** (8), 1476-
 374 1488. (doi:10.1111/jeb.12668).

375 18. Groothuis, T. G. G., Eising C. M., Dijkstra C. & Müller W. 2005 Balancing
 376 between costs and benefits of maternal hormone deposition in avian eggs. *Biol. Lett.* **1**
 377 (1), 78-81.

- 378 19. Tschirren, B., Richner H. & Schwabl H. 2004 Ectoparasite-modulated
379 deposition of maternal androgens in great tit eggs. *Proc. R. Soc. B* **271** (1546), 1371-
380 1375.
- 381 20. Tschirren, B., Sendecka J., Groothuis T. G. G., Gustafsson L. & Doligez B.
382 2009 Heritable variation in maternal yolk hormone transfer in a wild bird population
383 *Am. Nat.* **174** (4), 557-564.
- 384 21. Müller, W., Goerlich V. C., Vergauwen J., Groothuis T. G. G. & Eens M.
385 2012 Sources of variation in yolk hormone deposition: Consistency, inheritance and
386 developmental effects. *Gen. Comp. Endocrinol.* **175** (2), 337-343.
387 (doi:10.1016/j.ygcen.2011.11.034).
- 388 22. Okuliarová, M., Groothuis T. G. G., Skrobanek P. & Zeman M. 2011
389 Experimental evidence for genetic heritability of maternal hormone transfer to
390 offspring. *Am. Nat.* **177** (6), 824-834. (doi:10.1086/659996).
- 391 23. Pick, J. L., Hutter P. & Tschirren B. 2016 In search of genetic constraints
392 limiting the evolution of egg size: direct and correlated responses to artificial selection
393 on a prenatal maternal effector. *Heredity* **116**, 542–549. (doi:10.1038/hdy.2016.16).
- 394 24. Okuliarová, M., Škrobánek P. & Zeman M. 2009 Variability of yolk
395 testosterone concentrations during the reproductive cycle of Japanese quail. *Comp.*
396 *Biochem. Physiol. A* **154** (4), 530-534. (doi:10.1016/j.cbpa.2009.08.012).
- 397 25. Zeman, M., Kosutzky J. & Bobakova E. 1986 Testosterone concentration in
398 the seminal plasma of cocks. *Br. Poult. Sci.* **27** (2), 261-266.
399 (doi:10.1080/00071668608416879).
- 400 26. Pilz, K. M. & Smith H. G. 2004 Egg yolk androgen levels increase with
401 breeding density in the European Starling, *Sturnus vulgaris*. *Funct. Ecol.* **18** (1), 58-
402 66.

- 403 27. Mazuc, J., Bonneaud C., Chastel O. & Sorci G. 2003 Social environment
404 affects female and egg testosterone levels in the house sparrow (*Passer domesticus*).
405 *Ecol. Lett.* **6** (12), 1084-1090.
- 406 28. Verboven, N., Evans N. P., D'Alba L., Nager R. G., Blount J. D., Surai P. F. &
407 Monaghan P. 2005 Intra-specific interactions influence egg composition in the lesser
408 black-backed gull (*Larus fuscus*). *Behav. Ecol. Sociobiol.* **57** (4), 357-365.
409 (doi:10.1007/s00265-004-0862-x).
- 410 29. Gasparini, J., Boulinier T., Gill V. A., Gil D., Hatch S. A. & Roulin A. 2007
411 Food availability affects the maternal transfer of androgens and antibodies into eggs
412 of a colonial seabird. *J. Evol. Biol.* **20** (3), 874-880.
- 413 30. Giraudeau, M., Ziegler A. K., Pick J. L., Ducatez S., Canale C. I. & Tschirren
414 B. in press Interactive effects of yolk testosterone and carotenoid on pre-natal growth
415 and chick physiology. *Behav. Ecol.*
- 416 31. Burnham, K. P. & Anderson D. R. 2002 *Model selection and multimodel*
417 *inference: a practical information-theoretic approach*. 2nd ed ed. New York,
418 Springer-Verlag.
- 419 32. R Development Core Team. 2011 R: A language and environment for
420 statistical computing. <http://www.R-project.org>.
- 421 33. Bates, D., Maechler M. & Bolker B. 2011 lme4: Linear mixed-effects models
422 using S4 classes. (R package version 0.999375-42).
- 423 34. Ayers, K. L., Davidson N. M., Demiyah D., Roeszler K. N., Grützner F.,
424 Sinclair A. H., Oshlack A. & Smith C. A. 2013 RNA sequencing reveals sexually
425 dimorphic gene expression before gonadal differentiation in chicken and allows
426 comprehensive annotation of the W-chromosome. *Genome Biol.* **14** (3), doi:
427 10.1186/gb-2013-1114-1183-r1126. (doi:R26

428 10.1186/gb-2013-14-3-r26).

429 35. Smeds, L., Warmuth V., Bolivar P., Uebbing S., Burri R., Suh A., Nater A.,
430 Bureš S., Garamszegi L. Z., Hogner S., et al. 2015 Evolutionary analysis of the
431 female-specific avian W chromosome. *Nat. Commun.* **6**, doi:10.1038/ncomms8330.
432 (doi:733010.1038/ncomms8330).

433 36. Lemos, B., Branco A. T. & Hartl D. L. 2010 Epigenetic effects of
434 polymorphic Y chromosomes modulate chromatin components, immune response,
435 and sexual conflict. *Proc. Natl. Acad. Sci. U S A* **107** (36), 15826-15831.
436 (doi:10.1073/pnas.1010383107).

437 37. Dowling, D. K., Friberg U. & Lindell J. 2008 Evolutionary implications of
438 non-neutral mitochondrial genetic variation. *Trends Ecol. Evol.* **23** (10), 546-554.
439 (doi:10.1016/j.tree.2008.05.011).

440 38. Gemmell, N. J., Metcalf V. J. & Allendorf F. W. 2004 Mother's curse: the
441 effect of mtDNA on individual fitness and population viability. *Trends Ecol. Evol.* **19**
442 (5), 238-244. (doi:10.1016/j.tree.2004.02.002).

443 39. Novicic, Z. K., Immonen E., Jelic M., Andelkovic M., Stamenkovic-Radak M.
444 & Arnqvist G. 2015 Within-population genetic effects of mtDNA on metabolic rate in
445 *Drosophila subobscura*. *J. Evol. Biol.* **28** (2), 338-346. (doi:10.1111/jeb.12565).

446 40. Tschirren, B., Ziegler A. K., Canale C. I., Okuliarová M., Zeman M. &
447 Giraudeau M. in press High yolk testosterone transfer is associated with an increased
448 female metabolic rate. *Physiol. Biochem. Zool.*

449 41. Wright, A. E., Harrison P. W., Montgomery S. H., Pointer M. A. & Mank J. E.
450 2014 Independent stratum formation of the avian sex chromosomes reveals inter-
451 chromosomal gene conversion and predominance of purifying selection on the W
452 chromosome. *Evolution* **68** (11), 3281-3295. (doi:10.1111/evo.12493).

- 453 42. Backström, N., Ceplitis H., Berlin S. & Ellegren H. 2005 Gene conversion
454 drives the evolution of HINTW, an ampliconic gene on the female-specific avian W
455 chromosome. *Mol. Biol. Evol.* **22** (10), 1992-1999. (doi:10.1093/molbev/msi198).
- 456 43. Aiken, C. E. & Ozanne S. E. 2014 Transgenerational developmental
457 programming. *Hum. Reprod. Update* **20** (1), 63-75. (doi:10.1093/humupd/dmt043).
- 458 44. Danchin, E., Charmantier A., Champagne F. A., Mesoudi A., Pujol B. &
459 Blanchet S. 2011 Beyond DNA: integrating inclusive inheritance into an extended
460 theory of evolution. *Nat. Rev. Genet.* **12** (7), 475-486. (doi:10.1038/nrg3028).
- 461 45. Jablonka, E. & Raz G. 2009 Transgenerational epigenetic inheritance:
462 revalence, mechanisms, and implications for the study of heredity and evolution *Q.*
463 *Rev. Biol.* **84** (2), 131-176.
- 464 46. Müller, W., Deptuch K., Lopez-Rull I. & Gil D. 2007 Elevated yolk androgen
465 levels benefit offspring development in a between clutch context. *Behav. Ecol.* **18** (5),
466 929-936.
- 467 47. Pfannkuche, K. A., Gahr M., Weites I. M., Riedstra B., Wolf C. & Groothuis
468 T. G. G. 2011 Examining a pathway for hormone mediated maternal effects - Yolk
469 testosterone affects androgen receptor expression and endogenous testosterone
470 production in young chicks (*Gallus gallus domesticus*). *Gen. Comp. Endocrinol.* **172**
471 (3), 487-493. (doi:10.1016/j.ygcen.2011.04.014).
- 472 48. Rubolini, D., Martinelli R., von Engelhardt N., Romano M., Groothuis T. G.
473 G., Fasola M. & Saino N. 2007 Consequences of prenatal androgen exposure for the
474 reproductive performance of female pheasants (*Phasianus colchicus*). *Proc. R. Soc. B*
475 **274** (1606), 137-142. (doi:10.1098/rspb.2006.3696).

- 476 49. Müller, W., Groothuis T. G. G., Goerlich V. C. & Eens M. 2011 GnRH - a
477 Missing Link between Testosterone Concentrations in Yolk and Plasma and Its
478 Intergenerational Effects. *Plos One* **6** (7). (doi:e22675
479 10.1371/journal.pone.0022675).
- 480 50. Navara, K. J. 2013 Stress, hormones, and sex: How do we solve the puzzle of
481 sex ratio adjustment in birds? *Integ. Comp. Biol.* **53**, E153-E153.
- 482 51. Pilz, K. M., Adkins-Regan E. & Schwabl H. 2005 No sex difference in yolk
483 steroid concentrations of avian eggs at laying. *Biol. Lett.* **1** (3), 318-321.
484 (doi:10.1098/rsbl.2005.0321).
- 485 52. Muriel, J., Perez-Rodriguez L., Puerta M. & Gil D. 2015 Diverse dose-
486 response effects of yolk androgens on embryo development and nestling growth in a
487 wild passerine. *J. Exp. Biol.* **218** (14), 2241-2249. (doi:10.1242/jeb.118257).
- 488 53. Frésard, L., Morisson M., Brun J. M., Collin A., Pain B., Minvielle F. & Pitel
489 F. 2013 Epigenetics and phenotypic variability: some interesting insights from birds.
490 *Gen. Sel. Evol.* **45**, doi:10.1186/1297-9686-1145-1116. (doi:16
491 10.1186/1297-9686-45-16).
- 492 54. Moore, T. & Haig D. 1991 Genomic imprinting in mammaian development - a
493 parental tug-of-war *Trends Genet.* **7** (2), 45-49. (doi:10.1016/0168-9525(91)90040-w).
- 494 55. Frésard, L., Leroux S., Servin B., Gourichon D., Dehais P., San Cristobal M.,
495 Marsaud N., Vignoles F., Bed'hom B., Coville J. L., et al. 2014 Transcriptome-wide
496 investigation of genomic imprinting in chicken. *Nucleic Acids Res.* **42** (6), 3768-3782.
497 (doi:10.1093/nar/gkt1390).
- 498 56. Rice, W. R. 1984 Sex chromosomes and the evolution of sexual dimorphism
499 *Evolution* **38** (4), 735-742. (doi:10.2307/2408385).

500 57. Harano, T., Okada K., Nakayama S., Miyatake T. & Hosken D. J. 2010
501 Intralocus sexual conflict unresolved by sex-limited trait expression. *Curr. Biol.* **20**
502 (22), 2036-2039. (doi:10.1016/j.cub.2010.10.023).

503 58. Mills, S. C., Koskela E. & Mappes T. 2012 Intralocus sexual conflict for
504 fitness: sexually antagonistic alleles for testosterone. *Proc. R. Soc. B* **279** (1735),
505 1889-1895. (doi:10.1098/rspb.2011.2340).

506 59. Groothuis, T. G. G. & Schwabl H. 2008 Hormone-mediated maternal effects
507 in birds: mechanisms matter but what do we know of them? *Phil. Trans. R. Soc. B* **363**
508 (1497), 1647-1661. (doi:10.1098/rstb.2007.0007).

509 60. Schwabl, H. 1996 Environment modifies the testosterone levels of a female
510 bird and its eggs. *J. Exp. Zool.* **276** (2), 157-163.

511 61. Räsänen, K. & Kruuk L. E. B. 2007 Maternal effects and evolution at
512 ecological time-scales. *Funct. Ecol.* **21** (3), 408-421. (doi:10.1111/j.1365-
513 2435.2007.01246.x).

514 62. Hinde, C. A., Buchanan K. L. & Kilner R. M. 2009 Prenatal environmental
515 effects match offspring begging to parental provisioning. *Proc. R. Soc. B* **276** (1668),
516 2787-2794. (doi:10.1098/rspb.2009.0375).

517

518

Figure legends

Fig. 1. Resemblance in yolk testosterone deposition (log yolk T; pg / mg yolk) among family members. A) relationship between mothers and daughters; B) relationship between maternal grandmothers and granddaughters; C) relationship between paternal grandmothers and granddaughters.




Fig. 2. Effect of prenatal testosterone manipulation on the transfer of yolk testosterone to the eggs. Shown is the difference between the yolk testosterone concentration (log yolk T pg / mg yolk) in the eggs of females that have experienced an experimentally increased yolk testosterone level during their prenatal development (T) and females that developed in a control egg (C), and their mother. Means \pm SE are shown.

